

Cream and sterile gauze impregnated with a Hyaluronic Acid and Silver Sulfadiazine base

- Cream (tube x 100g)
- Cream (tube x 25g)
- Impregnated sterile gauze (box x 3 gauzes - 10 x 10cm)
- Impregnated sterile gauze (box x 10 gauzes - 10 x 10cm)

COMPOSITION

ialuset® Plus cream: Hyaluronic acid (sodium salt) 0.2%, Silver sulfadiazine 1%,

Other components: Decyl ester of oleic acid, Cetearyl alcohol and Sodium lauryl sulfate and Sodium cetearyl sulfate, Macrogol, Liquid sorbitol, Glycerol, Water.

ialuset® Plus impregnated sterile gauze: sterile gauze impregnated with cream of hyaluronic acid (sodium salt) 0.05% & silver sulfadiazine 1%,

Other components: Macrogol, Glycerol, Water.

ADMINISTRATION ROUTES AND POSOLOGY

Cream:

After cleaning and disinfecting the wound, apply 2-3 mm thick layer of cream directly onto the wound itself, or spread the cream on a sterile dressing and apply it onto the wound.

The dressing can be held in place with a suitable bandage.

It is recommended to clean the wound using physiological solution prior to each application. Apply every 24 hours from the budding phase until complete cicatrisation. The cream can be easily washed away using plain tap water.

Healing wounds and burns cicatrisation must be evaluated by a physician. The total duration treatment should not exceed one month.

Impregnated sterile gauze:

After cleaning and disinfecting the wound, remove the 2 protective foils from the gauze and apply it onto the wound. Then, the sterile impregnated gauze must be covered by a sterile dressing and a cotton pad.

The whole dressing has to be held in place with a suitable bandage.

In case of slight adhesion of the dressing to the wound, rinse thoroughly with physiological solution.

It is recommended to clean the wound using physiological solution prior to each application. Apply every 24 hours from the budding phase until complete cicatrisation.

Healing wounds and burns cicatrisation must be evaluated by a physician. The total duration treatment should not exceed one month.

UNDESIRABLE EFFECTS

Local hypersensitivity reactions, such as eczema have rarely been observed.

CONTRAINDICATIONS

Known sensitivity to sulfamides, general or topical (eczema through contact). Allergy to any of the components.

Due to insufficient data, the use of ialuset® Plus is counter-indicated in children less than 2 years of age.

ASK YOUR PHYSICIAN OR PHARMACIST FOR ADVICE AND TELL THEM ABOUT ANY ADVERSE REACTION YOU MAY EXPERIENCE AND WHICH IS NOT LISTED IN THIS LEAFLET.

PRECAUTIONS FOR USE

- Follow the proper methods for cleaning and disinfecting wounds, burns and the adjacent skin areas. The use of ialuset® Plus in association with an enzymatic detergent has not been evaluated.
- Avoid exposing to sun body areas on which ialuset® Plus is applied (due to risk of light sensitivity). Transient coloration of the wound bed may appear, which can be removed by simple washing.
- Patients with renal or hepatic deficiencies must avoid using the product on open wounds.
- Children: ialuset® Plus should be used with caution in children older than 2 years. The treatment duration should not exceed 15 days without medical advice.
- Keep out of reach of children.
- Do not swallow.

- Cream: Secure screw-cap carefully following each use. Do not directly apply the tube on the wound.
- Single use sterile impregnated gauze (Beta radiation): do not reuse to avoid any risk of contamination.
- Do not use if the packaging has suffered damage.
- Do not throw out the expired or unused products. Bring them to your pharmacist for disposal.
- Pregnancy and lactation: ialuset® Plus should only be used in pregnant and lactating women when strictly necessary.

IN CASE OF DOUBT, DO NOT HESITATE TO ASK YOUR PHYSICIAN OR PHARMACIST FOR ADVICE.

STORAGE

DO NOT USE AFTER THE EXPIRY DATE SHOWN ON THE PACKAGING.

Store at temperature below 30°C protected from light. Use within 6 months after opening (tube).

YEAR OF AUTHORIZATION OF THE CE MARK: 2000
LAST PATIENT INFORMATION LEAFLET REVIEW: Juin 2018



Warnings - See Precautions for Use



Sterilization by Radiation
 Only the impregnated gauze pads are sterile



Storage Temperature



Do not reuse - Single use only - Only the sterile impregnated gauze pads are for single use



Symbol for EC marking



Batch Number



Expiry Date/Use by



Laboratoires **Genevrier** - 280 rue de Goa - ZI les 3 Moulins
 Parc de Sophia Antipolis - 06901 Sophia Antipolis - France



Distributor: Name and Address of the Local Distributor